

The study by Bonello *et al*¹⁴, published in this issue of *Heart*, addresses prevention of myonecrosis using trimetazidine, an anti-ischaemic agent. In that study, 266 patients were randomly assigned to pretreatment with an oral loading dose of 60 mg trimetazidine or placebo. The group treated with trimetazidine had troponin values consistently lower than controls, and the total amount of post-PCI troponin release was reduced. This result was achieved despite a slight imbalance in baseline risk for periprocedural events against the trimetazidine group (in which patients were older, more frequently had diabetes mellitus and had a high body mass index and a high angina class). Because trimetazidine is devoid of haemodynamic effects and has overall excellent clinical tolerance, it is an attractive candidate as adjunctive treatment to PCI to decrease periprocedural myocardial injury. There are, however, limitations to this study: it is a single-centre, open study; in addition, although troponin release was reduced by trimetazidine, the frequency of troponin rise was not. Clearly, larger clinical studies powered on genuine clinical outcomes are warranted to confirm the findings of Bonello *et al* before such treatment can be adopted in clinical practice.

Authors' affiliations

Patrick Schiano, Hôpital d'Instruction des Armées du Val de Grâce, Paris, France

Philippe Gabriel Steg, Centre Hospitalier Bichat-Claude Bernard, Paris, France

Competing interests: None declared.

PGS has been a speaker and has served on advisory boards for Servier, the manufacturer of trimetazidine.

REFERENCES

- 1 Porto I, Selvanayagam JB, Van Gaal WJ, *et al*. Plaque volume and occurrence and location of periprocedural myocardial necrosis after percutaneous coronary intervention. Insights from delayed-enhancement magnetic resonance imaging, thrombolysis in myocardial infarction myocardial perfusion grade analysis, and intravascular ultrasound. *Circulation* 2006;**114**:662–9.
- 2 Miller LW, Garratt KN, Burritt MF, *et al*. Baseline troponin level: key to understanding the importance of post-PCI troponine elevations. *Eur Heart J* 2006;**27**:1061–9.
- 3 Cavallini C, Savonitto S, Violini R, on behalf the Italian "Atherosclerosis, thrombosis, and vascular biology" and "Society for invasive cardiology-GISE" investigators, *et al*. Impact of the elevation of biochemical markers of myocardial damage on long-term mortality after percutaneous coronary intervention: results of the CK-MB and PCI study. *Eur Heart J* 2005;**26**:1494–8.
- 4 Prasad A, Singh M, Lerman A, *et al*. Isolated elevation in troponin T after percutaneous coronary intervention is associated with higher long-term mortality. *J Am Coll Cardiol* 2006;**48**:1765–70.
- 5 Herrmann J. Peri-procedural myocardial injury: 2005 update. *Eur Heart J* 2005;**26**:2493–519.
- 6 Ioannidis JPA, Karvouni E, Katritsis G. Mortality risk conferred by small elevations of creatine kinase-MB isoenzyme after percutaneous coronary intervention. *J Am Coll Cardiol* 2003;**42**:1406–11.
- 7 Chew DP, Bhatt DL, Lincoff AM, *et al*. Clinical end point definitions after percutaneous coronary intervention and their relationship to late mortality: an assessment by attributable risk. *Heart* 2006;**92**:945–50.
- 8 Selvanayagam JB, Porto I, Channon K, *et al*. Troponin elevation after percutaneous coronary intervention directly represents the extent of irreversible myocardial injury. Insights from cardiovascular magnetic resonance imaging. *Circulation* 2005;**111**:1027–32.
- 9 Alpert JS, Thygesen K, on behalf of the Joint European Society of Cardiology/American College of Cardiology Committee, *et al*. Myocardial infarction redefined—a consensus document of the Joint European Society of Cardiology/American College of Cardiology committee for the redefinition of myocardial infarction. *Eur Heart J* 2000;**21**:1502–13.
- 10 Chen WH, Lee P, Ng W, *et al*. Aspirin resistance is associated with a high incidence of myonecrosis after non-urgent percutaneous coronary intervention despite clopidogrel pre-treatment. *J Am Coll Cardiol* 2004;**43**:1122–6.
- 11 Patti G, Colonna G, Pasceri V, *et al*. Randomized trial of high loading dose of clopidogrel for reduction of periprocedural myocardial infarction in patients undergoing coronary intervention. Results from the ARMYDA-2 (Antiplatelet therapy for Reduction of MYocardial Damage during Angioplasty) study. *Circulation* 2005;**111**:2099–106.
- 12 Anderson KM, Califf RM, Stone GW, *et al*. Long-term mortality benefit with abciximab in patients undergoing percutaneous coronary intervention. *J Am Coll Cardiol* 2001;**37**:2059–65.
- 13 Pasceri V, Patti G, Nusca A, on behalf of the ARMYDA investigators, *et al*. Randomized trial of atorvastatin for reduction of myocardial damage during coronary intervention. Results from the ARMYDA (Atorvastatin for Reduction of MYocardial Damage during Angioplasty) study. *Circulation* 2004;**110**:674–8.
- 14 Bonello L, Sbragia P, Amabile N, *et al*. Protective effect of an acute oral loading dose of trimetazidine on myocardial injury following percutaneous coronary intervention. *Heart* 2007;**93**:703–7.

bmjupdates+

bmjupdates+ is a unique and free alerting service, designed to keep you up to date with the medical literature that is truly important to your practice. bmjupdates+ will alert you to important new research and will provide you with the best new evidence concerning important advances in health care, tailored to your medical interests and time demands.

Where does the information come from?

bmjupdates+ applies an expert critical appraisal filter to over 100 top medical journals. A panel of over 2000 physicians find the few 'must read' studies for each area of clinical interest.

Sign up to receive your tailored email alerts, searching access and more...

www.bmjupdates.com